

Tennessee State University

Digital Scholarship @ Tennessee State University

Chemistry Faculty Research

Department of Chemistry

2009

Ethyl 1-acetyl-1H-indole-3-carboxylate

Tasneem Siddiquee

Tennessee State University

Shahid Islam

University of Wisconsin-Milwaukee

Dennis Bennett

University of Wisconsin-Milwaukee

Matthias Zeller

Youngstown State University

Mahmun Hossain

University of Wisconsin-Milwaukee

Follow this and additional works at: <https://digitalscholarship.tnstate.edu/chemistry-faculty>



Part of the [Chemistry Commons](#)

Recommended Citation

T. Siddiquee, S. Islam, D. Bennett, M. Zeller and M. Hossain "Ethyl 1-acetyl-1H-indole-3-carboxylate" *Acta Cryst.* (2009). E65, o1802-o1803 <https://doi.org/10.1107/S1600536809025379>

This Article is brought to you for free and open access by the Department of Chemistry at Digital Scholarship @ Tennessee State University. It has been accepted for inclusion in Chemistry Faculty Research by an authorized administrator of Digital Scholarship @ Tennessee State University. For more information, please contact XGE@Tnstate.edu.

Ethyl 1-acetyl-1*H*-indole-3-carboxylate

Tasneem Siddiquee,^{a*} Shahid Islam,^b Dennis Bennett,^b Matthias Zeller^c and Mahmun Hossain^b

^aDepartment of Chemistry, Boswell Science Complex, Tennessee State University, Nashville, 3500 John A Merritt Blvd, Nashville, TN 37209, USA, ^bDepartment of Chemistry and Biochemistry, University of Wisconsin-Milwaukee, 3210 N Cramer Street, Milwaukee, WI 53211, USA, and ^cYoungstown State University, Department of Chemistry, One University Plaza, Youngstown Ohio 44555-3663, USA
Correspondence e-mail: tsiddiqu@tnstate.edu


Received 11 June 2009; accepted 1 July 2009

Key indicators: single-crystal X-ray study; $T = 296\text{ K}$; mean $\sigma(\text{C}-\text{C}) = 0.002\text{ \AA}$; R factor = 0.042; wR factor = 0.120; data-to-parameter ratio = 13.1.

The title compound, $\text{C}_{13}\text{H}_{13}\text{NO}_3$, was synthesized by acetylation of ethyl 1*H*-indole-3-carboxylate. The aromatic ring system of the molecule is essentially planar, but the saturated ethyl group is also located within this plane and the overall r.m.s. deviation from planarity is only 0.034 \AA . Pairs of $\text{C}-\text{H}\cdots\text{O}$ interactions connect molecules into chains along the diagonal of the unit cell. Molecules also form weakly connected dimers *via* $\pi\cdots\pi$ stacking interactions of the indole rings with centroid–centroid separations of $3.571(1)\text{ \AA}$. $\text{C}-\text{H}\cdots\pi$ interactions between methylene and methyl groups and the indole and benzene ring complete the directional intermolecular interactions found in the crystal structure.

Related literature

For the biological properties of tryptophan derivatives, see: Ma *et al.* (2001); Zhou *et al.* (2006); Zhao, Smith *et al.* (2002); Zhao, Liao & Cook (2002). For synthetic procedures towards tryptophan-like compounds, see: Ager & Laneman (2004); Amir-Heidari *et al.* (2007); Carlier *et al.* (2002); Hengartner *et al.* (1979); Moriya *et al.* (1980). For the synthesis of 2-acetamido-3-ethoxy-3-oxopropanoic acid, see: Hellmann *et al.* (1958). For NMR data for the title compound, see: Reimann *et al.* (1990).

**Experimental***Crystal data*

$\text{C}_{13}\text{H}_{13}\text{NO}_3$	$\gamma = 114.28(1)^\circ$
$M_r = 231.24$	$V = 578.58(15)\text{ \AA}^3$
Triclinic, $P\bar{1}$	$Z = 2$
$a = 7.519(1)\text{ \AA}$	Mo $K\alpha$ radiation
$b = 8.479(1)\text{ \AA}$	$\mu = 0.10\text{ mm}^{-1}$
$c = 10.187(2)\text{ \AA}$	$T = 296\text{ K}$
$\alpha = 97.38(1)^\circ$	$0.51 \times 0.41 \times 0.20\text{ mm}$
$\beta = 95.78(2)^\circ$	

Data collection

Siemens P4 diffractometer	2027 independent reflections
Absorption correction: multi-scan [<i>XSCANS</i> (Siemens, 1996) and <i>XPREP</i> (Siemens, 1994)]	1696 reflections with $I > 2\sigma(I)$
$T_{\min} = 0.823$, $T_{\max} = 0.981$	$R_{\text{int}} = 0.019$
2536 measured reflections	3 standard reflections
	every 97 reflections
	intensity decay: <1%

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.042$	155 parameters
$wR(F^2) = 0.120$	H-atom parameters constrained
$S = 1.09$	$\Delta\rho_{\max} = 0.17\text{ e \AA}^{-3}$
2027 reflections	$\Delta\rho_{\min} = -0.18\text{ e \AA}^{-3}$

Table 1
Hydrogen-bond geometry (\AA , $^\circ$).

$D-\text{H}\cdots A$	$D-\text{H}$	$\text{H}\cdots A$	$D\cdots A$	$D-\text{H}\cdots A$
$\text{C}2-\text{H}2\cdots\text{O}3^{\text{i}}$	0.93	2.61	3.296 (2)	131
$\text{C}5-\text{H}5\cdots\text{O}1^{\text{ii}}$	0.93	2.64	3.273 (2)	125
$\text{C}12-\text{H}12\text{B}\cdots\text{Cg}1^{\text{iii}}$	0.96	2.95	3.618 (3)	127
$\text{C}13-\text{H}13\text{B}\cdots\text{Cg}2^{\text{iii}}$	0.96	2.78	3.587 (3)	142

Symmetry codes: (i) $-x + 1, -y + 2, -z + 1$; (ii) $-x, -y + 1, -z + 2$; (iii) $-x, -y + 2, -z + 2$. $\text{Cg}1$ is the centroid of the $\text{N}1,\text{C}1,\text{C}6-\text{C}8$ pyrrole ring and $\text{Cg}2$ is the centroid of the $\text{C}1-\text{C}6$ phenyl ring.

Data collection: *XSCANS* (Siemens, 1996); cell refinement: *XSCANS*; data reduction: *XSCANS*; program(s) used to solve structure: *XPREP* (Siemens 1994) and *SHELXTL* (Sheldrick, 2008); program(s) used to refine structure: *SHELXTL*; molecular graphics: *SHELXTL*; software used to prepare material for publication: *SHELXTL*.

TAS acknowledges the College of Arts and Science at TSU for release time.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: BH2233).

References

- Ager, D. J. & Laneman, S. (2004). *Asymmetric Catalysis on Industrial Scale: Challenges, Approaches and Solutions*, edited by H. U. Blaser & E. Schmidt, p. 30. Weinheim: WILEY-VCH Verlag GmbH & Co KGaA.
- Amir-Heidari, B., Thirlway, J. & Micklefield, J. (2007). *Org. Lett.* **9**, 1513–1516.
- Carlier, P. R., Lam, P. C.-H. & Wong, D. M. (2002). *J. Org. Chem.* **67**, 6256–6259.
- Hellmann, H., Teichmann, K. & Lingens, F. (1958). *Chem. Ber.* **91**, 2427–2431.
- Hengartner, U., Valentine, D. Jr., Johnson, K. K., Larscheid, M. E., Pigott, F., Scheidl, F., Scott, J. W., Sun, R. C., Townsend, J. M. & Williams, T. H. (1979). *J. Org. Chem.* **44**, 3741–3747.
- Ma, C., Liu, X., Li, X., Flippin-Anderson, J., Yu, S. & Cook, J. M. (2001). *J. Org. Chem.* **66**, 4525–4542.
- Moriya, T., Hagiwara, K. & Yoneda, N. (1980). *Chem. Pharm. Bull.* **28**, 1711–1721.
- Reimann, E., Hassler, T. & Lotter, H. (1990). *Arch. Pharm.* **323**, 255–258.

- Sheldrick, G. M. (2008). *Acta Cryst. A* **64**, 112–122.
- Siemens (1994). *XPREP*. Siemens Analytical X-ray Instruments Inc., Madison, Wisconsin, USA.
- Siemens (1996). *XSCANS*. Siemens Analytical X-ray Instruments Inc., Madison, Wisconsin, USA.
- Zhao, S., Liao, X. & Cook, J. M. (2002). *Org. Lett.* **4**, 687–690.
- Zhao, S., Smith, K. S., Deveau, A. M., Dieckhaus, C. M., Johnson, M. A., Macdonald, T. L. & Cook, J. M. (2002). *J. Med. Chem.* **45**, 1559–1562.
- Zhou, H., Liao, X., Yin, W., Ma, J. & Cook, J. M. (2006). *J. Org. Chem.* **71**, 251–259.

supporting information

Acta Cryst. (2009). E65, o1802–o1803 [doi:10.1107/S1600536809025379]

Ethyl 1-acetyl-1*H*-indole-3-carboxylate

Tasneem Siddiquee, Shahid Islam, Dennis Bennett, Matthias Zeller and Mahmun Hossain

S1. Comment

Indole substituted at 3-position leads to variety of compounds that are precursors to biologically active important alkaloids. One of the most important compounds of this type is tryptophan, which possesses anticancerous, antimalarial, antiamoebic, and antihypertensive activities (Ma *et al.*, 2001; Zhou *et al.*, 2006; Zhao, Smith *et al.* 2002; Zhao, Liao, & Cook, 2002). α,β -Dehydroaminoacid esters (*e.g.* **1**, Fig. 1) are precursors to synthesizing tryptophan derivatives, which upon hydrogenation yield optically active tryptophan and its analogues (Ager & Laneman, 2004).

α,β -Dehydroamino acid esters were also synthesized using Erlenmeyer condensation (Amir-Heidari *et al.*, 2007), Schmidt olefinations (Carlier *et al.*, 2002), condensation of indole aldehyde with acetyl amino malonic acid ester (Hengartner *et al.*, 1979), and Knoevenagel-type condensation (Moriya *et al.* 1980). One such effort to synthesize hydroxyl dehydrotryptophan (**3**) from indolester (**1**) using mono acid malonic ester (**2**) and acetic anhydride-pyridine mixture (Fig. 1) proved to be unsuccessful. The reaction resulted in 1*H*-indole-3-carboxylic acid-*N*-acetylethyl ester (**4**) instead. We rationalize that it is the electron withdrawing effect of the ester group which increases the acidity of the molecule. Consequently, in presence of a base, like pyridine, deprotonation and introduction of an acylium ion may occur. In this article we report the crystal structure of this compound.

The structure of the title compound is shown in Figure 2. The aromatic ring system of the molecule is essentially planar, but also the saturated ethyl group is located within this plane and the overall r.m.s. deviation from planarity is only 0.034 Å. Pairs of C—H···O interactions connect molecules into chains along the diagonal of the unit cell (Fig. 3). Molecules form weakly connected dimers *via* $\pi\cdots\pi$ stacking interactions of the indole rings with centroid to centroid distances of 3.571 (1) Å [symmetry operator for the second indole ring: (iii) 1 - x , 2 - y , 2 - z]. C—H··· π interactions between methylene and methyl groups and the indole and benzene ring complete the range of intermolecular interactions [C12—H12B···Cg1ⁱⁱⁱ = 2.95 Å, X—H···Cg1ⁱⁱⁱ = 127°, X···Cg1ⁱⁱⁱ = 3.618 (3) Å; C13—H13B···Cg2ⁱⁱⁱ = 2.78 Å, X—H···Cg2ⁱⁱⁱ = 142°, X···Cg2ⁱⁱⁱ = 3.587 (3) Å; Cg1 and Cg2 are the centroids of the indole and the benzene rings, respectively].

S2. Experimental

2-Acetamido-3-ethoxy-3-oxopropanoic acid (one of the starting materials) was prepared from acetyl amino malonic acid diethylester following the process developed by Hellmann *et al.* (1958). The title compound was prepared as follows: to a mixture of 0.37 g (1.97 mmol) of the indole ester ethyl 1*H*-indole-3-carboxylate, 1.1 g (5.9 mmol) of 2-acetamido-3-ethoxy-3-oxopropanoic acid, and 4.54 ml of pyridine was added at 288 K (15 °C) over 15 minutes 1.6 ml of acetic anhydride. The reaction mixture turned yellow and was stirred at 333 K (60 °C) for 3 h. An additional 0.18 g (0.9 mmol) of ethyl acetamido malonate was added and stirring was continued for 22 h. Ice (10 ml) was added, and the mixture was stirred for 2 h and then diluted with 20 ml of water. The resulting solution was extracted with EtOAc (2 × 20 ml), the combined organic layer was dried with anhydrous Na₂SO₄ and the solvent was removed under reduced pressure. 0.4 g (99%) of 1*H*-indole-3-carboxylic acid-*N*-acetyl ethyl ester was isolated. ¹HNMR CDCl₃, δ (p.p.m.): 8.70–8.50 (m, 1H and

2H), 7.60–7.30 (m, 2H), 4.45 (q, $J = 7$ Hz, OCH₂CH₃), 2.70 (s, COCH₃), 1.45 (t, $J = 7$ Hz, OCH₂CH₃). The NMR data agree with those reported previously (Reimann *et al.*, 1990). Crystals suitable for X-ray structural analysis were obtained by recrystallization from ethanol in a refrigerator.

S3. Refinement

All hydrogen atoms were added in calculated positions with a C—H bond distances of 0.97 (methylene), 0.93 (aromatic) and 0.96 Å (methyl). They were refined with isotropic displacement parameters U_{iso} of 1.5 (methyl) or 1.2 times U_{eq} (all others) of the adjacent carbon atom.






Figure 1

Synthesis of the title compound (**4**).

**Figure 2**

Thermal ellipsoid plot of the title compound with the atom labeling scheme. Displacement ellipsoids are shown at the 50% probability level and H atoms are shown as capped sticks.

**Figure 3**

Packing view of the title compound showing C—H···O interactions (blue lines).

Ethyl 1-acetyl-1*H*-indole-3-carboxylate*Crystal data*

$C_{13}H_{13}NO_3$
 $M_r = 231.24$
Triclinic, $P\bar{1}$
Hall symbol: -P 1
 $a = 7.519$ (1) Å
 $b = 8.479$ (1) Å
 $c = 10.187$ (2) Å
 $\alpha = 97.38$ (1)°
 $\beta = 95.78$ (2)°
 $\gamma = 114.28$ (1)°
 $V = 578.58$ (15) Å³

$Z = 2$
 $F(000) = 244$
 $D_x = 1.327$ Mg m⁻³
Mo $K\alpha$ radiation, $\lambda = 0.71073$ Å
Cell parameters from 23 reflections
 $\theta = 3.7$ –11.4°
 $\mu = 0.10$ mm⁻¹
 $T = 296$ K
Block, colourless
 $0.51 \times 0.41 \times 0.20$ mm

Data collection

Siemens P4
diffractometer
Radiation source: fine-focus sealed tube
Graphite monochromator
 $2\theta/\omega$ scans
Absorption correction: multi-scan
[XSCANS (Siemens 1996) and XPREP
(Siemens, 1994)]
 $T_{\min} = 0.823$, $T_{\max} = 0.981$
2536 measured reflections

2027 independent reflections
1696 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.019$
 $\theta_{\max} = 25.0$ °, $\theta_{\min} = 2.1$ °
 $h = -8 \rightarrow 1$
 $k = -9 \rightarrow 9$
 $l = -12 \rightarrow 12$
3 standard reflections every 97 reflections
intensity decay: <1%

Refinement

Refinement on F^2
Least-squares matrix: full
 $R[F^2 > 2\sigma(F^2)] = 0.042$
 $wR(F^2) = 0.120$
 $S = 1.09$
2027 reflections
155 parameters
0 restraints
0 constraints
Primary atom site location: structure-invariant
direct methods
Secondary atom site location: difference Fourier
map

Hydrogen site location: inferred from
neighbouring sites
H-atom parameters constrained
 $w = 1/[\sigma^2(F_o^2) + (0.0647P)^2 + 0.0738P]$
where $P = (F_o^2 + 2F_c^2)/3$
 $(\Delta/\sigma)_{\max} = 0.001$
 $\Delta\rho_{\max} = 0.17$ e Å⁻³
 $\Delta\rho_{\min} = -0.18$ e Å⁻³
Extinction correction: SHELXTL (Bruker, 2003;
Sheldrick, 2008),
 $F_c^* = kFc[1 + 0.001xFc^2\lambda^3/\sin(2\theta)]^{-1/4}$
Extinction coefficient: 0.103 (12)

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (Å²)

	<i>x</i>	<i>y</i>	<i>z</i>	$U_{\text{iso}}^*/U_{\text{eq}}$
C1	0.3179 (2)	0.9158 (2)	0.76860 (15)	0.0474 (4)
C2	0.3276 (2)	0.8319 (2)	0.64495 (16)	0.0560 (4)
H2	0.3926	0.8952	0.5826	0.067*
C3	0.2364 (3)	0.6502 (2)	0.61896 (18)	0.0641 (5)
H3	0.2402	0.5898	0.5371	0.077*
C4	0.1394 (3)	0.5554 (2)	0.71141 (19)	0.0658 (5)
H4	0.0804	0.4330	0.6907	0.079*
C5	0.1286 (3)	0.6392 (2)	0.83389 (17)	0.0572 (4)

H5	0.0629	0.5747	0.8955	0.069*
C6	0.2186 (2)	0.8227 (2)	0.86285 (15)	0.0477 (4)
C7	0.2359 (2)	0.9517 (2)	0.97738 (15)	0.0477 (4)
C8	0.3421 (2)	1.1130 (2)	0.95024 (15)	0.0491 (4)
H8	0.3747	1.2189	1.0077	0.059*
C9	0.1539 (2)	0.9133 (2)	1.10037 (16)	0.0526 (4)
C10	0.5038 (3)	1.2371 (2)	0.76197 (16)	0.0563 (4)
C11	0.5648 (3)	1.4194 (3)	0.8383 (2)	0.0756 (6)
H11A	0.6392	1.4336	0.9248	0.113*
H11B	0.4491	1.4377	0.8497	0.113*
H11C	0.6452	1.5037	0.7894	0.113*
C12	0.1351 (3)	1.0333 (3)	1.31867 (17)	0.0613 (5)
H12A	0.1950	0.9683	1.3634	0.074*
H12B	-0.0076	0.9667	1.3045	0.074*
C13	0.1950 (3)	1.2107 (3)	1.40273 (18)	0.0724 (5)
H13A	0.1508	1.1970	1.4875	0.109*
H13B	0.1361	1.2743	1.3572	0.109*
H13C	0.3366	1.2747	1.4175	0.109*
N1	0.39555 (19)	1.09788 (17)	0.82406 (12)	0.0489 (4)
O1	0.0559 (3)	0.76795 (18)	1.11828 (14)	0.0845 (5)
O2	0.20161 (17)	1.05915 (15)	1.19112 (11)	0.0559 (3)
O3	0.5434 (2)	1.20764 (19)	0.65295 (13)	0.0814 (5)

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
C1	0.0485 (8)	0.0506 (9)	0.0420 (8)	0.0226 (7)	0.0046 (6)	0.0028 (6)
C2	0.0594 (10)	0.0612 (10)	0.0452 (9)	0.0262 (8)	0.0100 (7)	0.0006 (7)
C3	0.0707 (11)	0.0617 (11)	0.0531 (10)	0.0282 (9)	0.0083 (8)	-0.0092 (8)
C4	0.0744 (11)	0.0493 (10)	0.0660 (11)	0.0245 (9)	0.0072 (9)	-0.0049 (8)
C5	0.0631 (10)	0.0494 (9)	0.0552 (10)	0.0216 (8)	0.0084 (8)	0.0070 (7)
C6	0.0492 (8)	0.0497 (8)	0.0437 (8)	0.0231 (7)	0.0043 (6)	0.0036 (7)
C7	0.0521 (8)	0.0502 (9)	0.0419 (8)	0.0239 (7)	0.0077 (6)	0.0062 (7)
C8	0.0569 (9)	0.0503 (9)	0.0405 (8)	0.0245 (7)	0.0102 (6)	0.0032 (6)
C9	0.0617 (9)	0.0541 (10)	0.0467 (9)	0.0283 (8)	0.0122 (7)	0.0108 (7)
C10	0.0656 (10)	0.0570 (10)	0.0478 (9)	0.0254 (8)	0.0155 (7)	0.0131 (7)
C11	0.0980 (15)	0.0525 (11)	0.0726 (12)	0.0247 (10)	0.0284 (11)	0.0146 (9)
C12	0.0718 (11)	0.0789 (12)	0.0442 (9)	0.0395 (10)	0.0208 (8)	0.0160 (8)
C13	0.0812 (13)	0.0925 (14)	0.0491 (10)	0.0451 (11)	0.0156 (9)	0.0023 (9)
N1	0.0562 (8)	0.0484 (7)	0.0406 (7)	0.0215 (6)	0.0105 (5)	0.0044 (5)
O1	0.1286 (12)	0.0546 (8)	0.0692 (9)	0.0304 (8)	0.0424 (8)	0.0197 (6)
O2	0.0660 (7)	0.0586 (7)	0.0430 (6)	0.0256 (6)	0.0172 (5)	0.0069 (5)
O3	0.1158 (11)	0.0731 (9)	0.0556 (8)	0.0346 (8)	0.0377 (7)	0.0162 (7)

Geometric parameters (\AA , $^\circ$)

C1—C2	1.389 (2)	C9—O1	1.197 (2)
C1—C6	1.399 (2)	C9—O2	1.3367 (19)

C1—N1	1.4186 (19)	C10—O3	1.201 (2)
C2—C3	1.379 (2)	C10—N1	1.400 (2)
C2—H2	0.9300	C10—C11	1.497 (3)
C3—C4	1.384 (3)	C11—H11A	0.9600
C3—H3	0.9300	C11—H11B	0.9600
C4—C5	1.381 (2)	C11—H11C	0.9600
C4—H4	0.9300	C12—O2	1.449 (2)
C5—C6	1.393 (2)	C12—C13	1.494 (3)
C5—H5	0.9300	C12—H12A	0.9700
C6—C7	1.449 (2)	C12—H12B	0.9700
C7—C8	1.352 (2)	C13—H13A	0.9600
C7—C9	1.467 (2)	C13—H13B	0.9600
C8—N1	1.391 (2)	C13—H13C	0.9600
C8—H8	0.9300		
C2—C1—C6	122.32 (15)	O2—C9—C7	112.43 (14)
C2—C1—N1	130.20 (15)	O3—C10—N1	120.26 (16)
C6—C1—N1	107.48 (13)	O3—C10—C11	123.11 (16)
C3—C2—C1	116.89 (17)	N1—C10—C11	116.63 (15)
C3—C2—H2	121.6	C10—C11—H11A	109.5
C1—C2—H2	121.6	C10—C11—H11B	109.5
C2—C3—C4	121.75 (17)	H11A—C11—H11B	109.5
C2—C3—H3	119.1	C10—C11—H11C	109.5
C4—C3—H3	119.1	H11A—C11—H11C	109.5
C5—C4—C3	121.27 (17)	H11B—C11—H11C	109.5
C5—C4—H4	119.4	O2—C12—C13	107.88 (15)
C3—C4—H4	119.4	O2—C12—H12A	110.1
C4—C5—C6	118.33 (17)	C13—C12—H12A	110.1
C4—C5—H5	120.8	O2—C12—H12B	110.1
C6—C5—H5	120.8	C13—C12—H12B	110.1
C5—C6—C1	119.44 (14)	H12A—C12—H12B	108.4
C5—C6—C7	133.47 (15)	C12—C13—H13A	109.5
C1—C6—C7	107.10 (14)	C12—C13—H13B	109.5
C8—C7—C6	107.50 (14)	H13A—C13—H13B	109.5
C8—C7—C9	126.52 (15)	C12—C13—H13C	109.5
C6—C7—C9	125.98 (15)	H13A—C13—H13C	109.5
C7—C8—N1	110.33 (14)	H13B—C13—H13C	109.5
C7—C8—H8	124.8	C8—N1—C10	126.27 (14)
N1—C8—H8	124.8	C8—N1—C1	107.60 (13)
O1—C9—O2	123.51 (15)	C10—N1—C1	126.12 (13)
O1—C9—C7	124.06 (16)	C9—O2—C12	116.25 (13)
C6—C1—C2—C3	-0.9 (2)	C8—C7—C9—O1	177.82 (17)
N1—C1—C2—C3	-179.95 (15)	C6—C7—C9—O1	-2.6 (3)
C1—C2—C3—C4	0.1 (3)	C8—C7—C9—O2	-2.5 (2)
C2—C3—C4—C5	0.5 (3)	C6—C7—C9—O2	177.09 (13)
C3—C4—C5—C6	-0.2 (3)	C7—C8—N1—C10	179.14 (15)
C4—C5—C6—C1	-0.6 (2)	C7—C8—N1—C1	0.04 (17)

C4—C5—C6—C7	179.54 (17)	O3—C10—N1—C8	179.02 (16)
C2—C1—C6—C5	1.2 (2)	C11—C10—N1—C8	-1.0 (3)
N1—C1—C6—C5	-179.57 (13)	O3—C10—N1—C1	-2.1 (3)
C2—C1—C6—C7	-178.95 (14)	C11—C10—N1—C1	177.96 (15)
N1—C1—C6—C7	0.31 (16)	C2—C1—N1—C8	178.95 (16)
C5—C6—C7—C8	179.58 (17)	C6—C1—N1—C8	-0.23 (16)
C1—C6—C7—C8	-0.29 (17)	C2—C1—N1—C10	-0.1 (3)
C5—C6—C7—C9	0.0 (3)	C6—C1—N1—C10	-179.32 (15)
C1—C6—C7—C9	-179.90 (14)	O1—C9—O2—C12	2.7 (2)
C6—C7—C8—N1	0.15 (18)	C7—C9—O2—C12	-177.03 (13)
C9—C7—C8—N1	179.76 (14)	C13—C12—O2—C9	-177.54 (14)

Hydrogen-bond geometry (Å, °)

D—H···A	D—H	H···A	D···A	D—H···A
C2—H2···O3 ⁱ	0.93	2.61	3.296 (2)	131
C5—H5···O1 ⁱⁱ	0.93	2.64	3.273 (2)	125
C12—H12B···Cg1 ⁱⁱⁱ	0.96	2.95	3.618 (3)	127
C13—H13B···Cg2 ⁱⁱⁱ	0.96	2.78	3.587 (3)	142

Symmetry codes: (i) $-x+1, -y+2, -z+1$; (ii) $-x, -y+1, -z+2$; (iii) $-x, -y+2, -z+2$.