

Tennessee State University

Digital Scholarship @ Tennessee State University

Agricultural and Environmental Sciences
Faculty Research

Department of Agricultural and Environmental
Sciences

3-7-2017

Effect of Organic and Inorganic Fertilizers on Soil Properties and the Growth, Yield and Quality of Tomato in Mymensingh, Bangladesh

M. Ashraful Islam

Bangladesh Agriculture University

Sumiya Islam

Bangladesh Agriculture University

Ayasha Akter

Bangladesh Agriculture University

Md Habibur Rahman

Bangladesh Agriculture University

Dilip Nandwani

Tennessee State University

Follow this and additional works at: <https://digitalscholarship.tnstate.edu/agricultural-and-environmental-sciences-faculty>



Part of the [Agriculture Commons](#), and the [Plant Sciences Commons](#)

Recommended Citation

Islam, M.A.; Islam, S.; Akter, A.; Rahman, M.H.; Nandwani, D. Effect of Organic and Inorganic Fertilizers on Soil Properties and the Growth, Yield and Quality of Tomato in Mymensingh, Bangladesh. *Agriculture* 2017, 7, 18. <https://doi.org/10.3390/agriculture7030018>

This Article is brought to you for free and open access by the Department of Agricultural and Environmental Sciences at Digital Scholarship @ Tennessee State University. It has been accepted for inclusion in Agricultural and Environmental Sciences Faculty Research by an authorized administrator of Digital Scholarship @ Tennessee State University. For more information, please contact XGE@Tnstate.edu.

Article

Effect of Organic and Inorganic Fertilizers on Soil Properties and the Growth, Yield and Quality of Tomato in Mymensingh, Bangladesh

M. Ashraful Islam ¹, Sumiya Islam ¹, Ayasha Akter ¹, Md Habibur Rahman ¹
and Dilip Nandwani ^{2,*}

¹ Department of Horticulture, Bangladesh Agriculture University (BAU), Mymensingh 2202, Bangladesh; ashrafulmi@bau.edu.bd (M.A.I.); sumyaislam1990@gmail.com (S.I.); aakter.akhy@gmail.com (A.A.); mhrahmand3@yahoo.com (M.H.R.)

² Department of Agriculture and Environmental Sciences, College of Agriculture, Human and Natural Sciences, Tennessee State University, Nashville, TN 37209, USA

* Correspondence: dnandwan@tnstate.edu; Tel.: +1-615-963-1897

Academic Editor: Guodong Liu

Received: 30 December 2016; Accepted: 2 March 2017; Published: 7 March 2017

Abstract: Field trials were conducted on tomato for yield and quality of fruits using different types of organic and inorganic fertilizers at the horticulture farm of Bangladesh Agricultural University (BAU), Mymensingh. Fertilizer treatments were tested on two varieties of tomato ca. Roma VF and BARI 15. The fertilization treatments were T₁, vermicompost (12 t/ha); T₂, compost (10 t/ha); T₃, integrated plant nutrient system (IPNS) or mixed fertilizers (organic 2/3 part and inorganic 1/3 part); T₄, inorganic fertilizers; and a control (T₅). Results showed growth and yield (20.8 t/ha) in tomato were higher in the IPNS treatment. A higher number of fruits per plant (73.7) and plant height (73.5 cm) were obtained from mixed fertilizers (organic 2/3 + inorganic 1/3) or IPNS (integrated plant nutrient system) in Roma VF than other treatments. Fruit yield and diameter were found statistically significant. No significant difference was observed in the quality (total soluble solids) of tomato fruits in both varieties' response to the treatments. The electrical conductivity and pH of the soil were improved by the application of organic manure.

Keywords: compost; nutrient management; production; sustainable; vegetables

1. Introduction

The organic production system aims at supporting and sustaining healthy ecosystems, soil, farmers, food production, the community, and the economy. Reduction and elimination of the adverse effects of synthetic fertilizers and pesticides on human health and the environment is a strong indicator that organic agriculture is gaining worldwide attention [1,2]. Organic fertilizers are environmentally friendly, since they are from organic sources [3]. The current global scenario firmly emphasizes the need to adopt eco-friendly agricultural practices for sustainable food production. The cost of inorganic fertilizers is increasing enormously, to the extent that they are out of reach for small and marginal farmers. The organic fertilizers provide nutritional requirements, suppress plant pest populations, and increase the yield and quality of agricultural crops in ways similar to inorganic fertilizers [4–10]. Solaiman and Rabbani [11] reported results and their findings on the application of a combination of cow dung and a half dose of inorganic fertilizer. Bio fertilizers such as *Trichoderma* enriched with inorganic fertilizers play a significant role in the growth and yield of crops, e.g., mustard and tomato, and have the potential to reduce 50% of the cost of inorganic fertilizers [12]. Microorganisms or plant growth promoting rhizobacteria (*Bacillus* spp) inoculant help to reduce the application of inorganic

fertilizers and contribute to improving soil fertility and reducing a negative environmental impact [13]. Organic fertilizers contain macro-nutrients, essential micro-nutrients, vitamins, growth-promoting indole acetic acid (IAA), gibberellic acid (GA) and beneficial microorganisms [14]. The sustainability of conventional agriculture in Bangladesh is under threat from the continuous degradation of land and water resources and from declining yields due to the indiscriminate use of agro-chemicals. Soil quality has been affected by excessive applications of chemical farming in Bangladesh [15]. Tomato (*Solanum lycopersicum*) is one of the important vegetable crops grown throughout the world and ranks next to the potato and sweet potato in terms of area, but ranks first as a processing crop [16]. The cultivated area under tomato was 18.8 thousand hectares with a total production in Bangladesh of 251 thousands metric tons [17]. Tomato is grown in the summer and winter seasons in Bangladesh; however, production varies in various regions due to varieties, seasons, climatic conditions, planting times, management practices and soil properties [18,19]. Ahammad et al. [20] found that planting time and varietal selection influenced tomato production from 4.51 to 55.91 t/ha. The objective of this research project was to evaluate the effect of organic and inorganic fertilizers combined on the growth, yield and quality of tomato and on soil properties.

2. Materials and Methods

An open-field experiment was conducted at the Horticulture Farm (24°26' and 24°54' N latitude and 90°15' and 90°30' E longitude), Bangladesh Agricultural University (BAU), Mymensingh in Fall 2014 to Spring 2015.

Two factors experiment was laid out in the Randomized Complete Block Design (RCBD) with three replications. The double factor experiment consisted of 5 treatments (T) is presented in Table 1:

Table 1. Fertilizer treatments and doses.

Treatments (T)	Dose/ha
Vermicompost (T ₁)	12 ton
Compost (T ₂)	10 ton
Integrated plant nutrient system (IPNS)/Mixed fertilizers (T ₃)	(organic 2/3 + inorganic 1/3); 8 ton vermicompost + N = 40 kg, P = 14 kg, K = 24 kg, S = 5 kg, Zn = 0.7 kg, B = 700 g N = 120 kg, P = 40 kg, K = 70 kg, S = 15 kg, Zn = 2 kg, B = 1 kg
Inorganic fertilizers (T ₄)	
Control (T ₅)	No fertilizers and manures

Two varieties of tomato ca Roma VF (V₁) and BARI tomato 15 (V₂) were selected for the study. Soil samples were collected from the experimental field before and after research trial and analyzed at the Agrivarsity Humboldt Soil Testing Laboratory (Table 2). Seeds were sown in nursery beds (3 m × 1 m), one seedbed/variety. Weeding, mulching and watering in seedbeds were done as needed. Seedlings germinated within a week and transplanted (25 days) in the field. Plant spacing kept at 80 cm × 60 cm in the field plots (2.8 m × 1.8 m) and watered. Vermicompost and compost was applied during land preparation. Urea, MoP, TSP, Borax and Zn fertilizers were applied at recommended doses in the inorganic plots (Table 1). Nitrogen and potassium was applied in two equal splits at 15 and 35 days after transplanting (DAT). The research plot was irrigated after fertilizer application. Botanical pesticides (neem extract) were used to control the insect pests and diseases except the T₄ experimental plot where inorganic fertilizers and fungicide (Dithane M-45 @2 g/L fortnightly during the early vegetative stage) were applied. Staking was done with bamboo sticks to support tomato plants. Data were collected from five plants of nine planted in each plot (one replication).

Table 2. Soil nutrients before planting and after harvesting of tomato.

	pH	EC ($\mu\text{c}/\text{cm}$)	O.M (%)	N (%)	P (ppm)	K (ppm)	S (ppm)	Clay (%)	Slit (%)	Sand (%)	Soil Texture
Before planting	6.43	72.54	2.55	0.11	15.38	47.13	13.18	6.15	62.62	35.40	Silt loam
After harvesting	6.68	98.54	2.60	0.11	15.54	50.13	13.18	6.15	62.62	35.40	Silt loam

Plant height (15, 35 and 45 DAT) was recorded (cm) from the ground to the tip of stem. Data on number of fruits, flower clusters and number of fruit clusters per plant were recorded. Twenty fruits per plot and replication were measured for fruit length (cm) and fruit diameter (cm) by caliper. Total seven harvest recorded from 20 January to 15 February. Fruits were harvested at the maturing index (when fruits become 10% red color) and stored at the horticulture research laboratory, and total soluble solid (TSS) recorded from the marketable fruits. Mean comparisons between treatments were performed by the least significant difference (LSD) test. All statistical calculations were performed using F-variance test, statistical significance was indicated at 5% and 1% level of probability [21].

3. Results and Discussion

The analysis of variance results showed that the effect of mixed fertilizers on tomato yield was significant ($p < 0.01$). Higher yields were obtained from mixed fertilizer treatments compared to the control (no fertilizer application). The application of T₃ (mixed fertilizers, organic 2/3 + inorganic 1/3) produced the highest number of flower clusters (31.2), fruit clusters (24.9), fruit yield (15.3 t/ha) and plant height (71.6 cm). T₅ (control or no fertilizer) had the lowest number of flower clusters per plant (27.22), number of fruit clusters per plant (20.1 cm), yield (9.4 t/ha) and plant height (62.1 t/ha).

The plant height was significantly different in different fertilizer treatments between varieties. The maximum plant height (73.5 cm) was recorded in the IPNS (organic 2/3 + inorganic 1/3) application (Figure 1) in variety 1 (Roma VF) (Figure 2). IPNS treatment produced a 38.5% higher yield compared to the control. Similarly, 7.3%, 8.2%, 21.8% higher yields were recorded in the vermicompost, compost and chemical fertilizer applications respectively, compared to the control (Table 3). The results obtained concur with earlier findings reported by Patil et al. [22] and Reddy et al. [23]. The mixed fertilizer treatment was the best for tomato production, which could be interpreted as the release of nutrients from organic and inorganic fertilizers at different times. Fertilizer treatment (mixed or IPNS) produced a higher yield compared to organic fertilizers and inorganic fertilizers alone in cabbage [24] and in eggplant [25].




Figure 1. Effect of fertilizer treatments on plant height on different days after transplanting (DAT). Vertical bars represent Least Significant difference (LSD) at 1% level of probability. T₁ = Vermicompost, T₂ = Compost, T₃ = Integrated plant nutrient system (IPNS) (organic 2/3 + inorganic 1/3), T₄ = Inorganic fertilizers, T₅ = Control.




Figure 2. Plant height of two tomato varieties on different days after transplanting (DAT). Vertical bar represents LSD at 1% level of probability (V₁ = Roma VF and V₂ = BARI tomato 15).

Table 3. Effect of various fertilizer treatments on tomato growth, fruit characteristics and yield.

Treatment	No of Fruits per Plant	No of Flower Clusters per Plant	No of Fruit Clusters per Plant	Fruit Length (cm)	Fruit Diameter (cm)	Yield (t/ha)
T ₁	47.23	30.22	18.50	5.44	4.62	10.20
T ₂	58.28	27.22	24.11	5.21	4.61	10.30
T ₃	61.06	31.28	24.95	5.45	4.84	15.39
T ₄	64.12	27.89	22.39	5.32	4.62	12.09
T ₅	48.61	27.61	20.17	5.20	4.64	9.46
LSD _{0.05}	2.311	0.933	0.674	0.094	0.054	0.467
LSD _{0.01}	3.166	1.278	0.924	0.129	0.074	0.639
Level of significance	**	**	**	**	**	**

** indicate at the 1% level of significance.

Fertilizers and variety treatments influenced the number of fruits per plant. A higher number of fruits (73.7) was recorded in var. Roma VF (V₁T₃) with the mixed fertilizer treatment (organic 2/3 + inorganic 1/3), while the lowest number (41.3) was obtained in the control (no fertilizer) of the BARI tomato 15 (V₂T₅). Plant height of 61.1 cm was recorded in variety BARI tomato 15 in the control group (no fertilizer, V₂T₅) (Table 4). Both Roma VF (V₁) and BARI 15 (V₂) varieties showed enhanced performance with IPNS or mixed fertilizer applications. The results indicate that mixed fertilizers produced improved yield in tomato varieties. A high yield per hectare (15.3 t/ha) was obtained in T₃ (Mixed, organic 2/3 + inorganic 1/3), 12.1 t/ha in inorganic fertilizer (T₄), 10.2 t/ha in vermicompost (T₁) and 9.4 t/ha in compost (T₂) applications. Roma VF had significantly higher yields (12.8 t/ha) than BARI tomato 15 (10.1 t/ha). The maximum fruit yield (20.8 t/ha) was obtained from the mixed fertilizers (organic 2/3 + inorganic 1/3) with Roma VF (V₁T₃) and the minimum (8.4 t/ha) from the control (no fertilizer) with BARI tomato 15 (V₂T₅) (Table 5). The fruit length and diameter were higher under the IPNS treatment. Both varieties produced higher numbers of flower clusters and fruit clusters per plant. These findings trended significantly towards the higher production of tomato under the IPNS.

Roma VF produced a significantly higher yield (12.8 t/ha) than BARI tomato 15 (10.1 t/ha). A high fruit yield (20.8 t/ha) was obtained in the mixed fertilizer application (organic 2/3 + inorganic 1/3) for Roma VF (V₁T₃) and the lowest yield (8.4 t/ha) was obtained in the control (no fertilizer) with

BARI tomato 15 (V₂T₅). A significant difference in the fruit length and diameter was recorded in the two varieties. Flower and fruit clusters per plant were higher in Roma VF. The higher number of flower and fruit clusters per plant in both varieties was observed in the IPNS application (T₃). Combined effect of fertilizers and variety treatment observed higher yield in Roma VF, and the highest production under IPNS treatment of both varieties.

The improved yield in tomato in the combination of cow dung and inorganic fertilizer has been reported [11] in the literature. The enhanced performance of integrated nutrient management practices has been reported in vegetables such as brinjal, cabbage and mustard [12,24,25]. The application of the mixed fertilizer strategy and bio fertilizer *Trichoderma*-enriched fertilizer or the inoculation of rhizobacteria is gaining popularity in Bangladesh [12,13]. It has been found that microorganisms and plant growth promoting rhizobacteria (PGPR) *Bacillus* spp. enhance nutrient availability, nutrient use efficiency and increase the growth and yield of plants. Plant growth may be influenced by some PGPR through synthesizing plant hormones or facilitating the uptake of nutrients from the soil by different direct mechanisms, e.g., atmospheric nitrogen (N) fixation, solubilization of phosphorus (P), and synthesis of siderophores for iron sequestration, making nutrients available to plants [26,27].

Table 4. Combined effects of fertilizers and varieties on yield, plant height and fruits measurements.

Treatment Combination ¹	Plant Height (cm) at DAT ²			No of Fruits per Plant	No of Flower Cluster per Plant	No of Fruit Clusters per Plant	Fruit Length (cm)	Fruit Diameter (cm)	Yield (t/ha)
	15	30	45						
V ₁ T ₁	33.61	43.75	63.95	47.89	33.11	17.56	5.47	4.68	11.16
V ₁ T ₂	32.07	42.10	62.59	70.44	28.22	25.44	5.21	4.62	10.12
V ₁ T ₃	41.61	51.27	73.53	73.78	36.11	30.78	5.58	4.89	20.84
V ₁ T ₄	35.97	45.10	73.38	60.67	28.89	22.89	5.29	4.69	11.71
V ₁ T ₅	31.21	41.68	63.31	55.89	30.22	23.22	5.32	4.74	10.45
V ₂ T ₁	35.05	45.32	62.82	46.56	27.33	19.44	5.40	4.56	9.23
V ₂ T ₂	35.85	44.23	63.53	46.11	26.22	22.78	5.20	4.59	10.49
V ₂ T ₃	30.23	44.72	64.87	48.33	26.44	19.11	5.32	4.79	9.94
V ₂ T ₄	37.50	48.26	69.81	67.56	26.89	21.89	5.35	4.54	12.48
V ₂ T ₅	30.06	39.09	61.07	41.33	25.00	17.11	5.08	4.53	8.47
LSD _{0.05}	1.247	0.932	2.189	3.268	1.320	0.954	5.47	4.68	11.16
LSD _{0.01}	1.708	1.276	2.999	4.477	1.808	1.306	5.21	4.62	10.12
Level of significance ³	**	**	**	**	**	**	**	**	**

¹ T₁ = Vermicompost, T₂ = Compost, T₃ = Integrated plant nutrient system (IPNS) (organic 2/3 + inorganic 1/3), T₄ = Inorganic fertilizers, T₅ = Control, V₁ = Roma VF, V₂ = BARI tomato 15, ² DAT: Days after transplanting, ³,** indicate at the 1% level of significance.

Table 5. Fruit characteristics and yield in tomato varieties.

Variety	No of Fruits per Plant	No of Flower Clusters per Plant	No of Fruit Clusters per Plant	Fruit Length (cm)	Fruit Diameter (cm)	Yield (t/ha)
V ₁	61.73	31.31	23.98	5.37	4.72	12.85
V ₂	49.98	26.38	20.07	5.27	4.60	10.12
LSD _{0.05}	1.462	0.590	0.426	0.059	0.034	0.295
LSD _{0.01}	2.002	0.809	0.584	0.081	0.047	0.404
Level of significance	**	**	**	**	**	**

** indicate at the 1% level of significance, V₁ = Roma VF, V₂ = BARI tomato 15.

The total soluble solid (TSS) of Roma VF (V₁) and BARI 15 (V₂) was 9 and 10, respectively. Fertilizer treatments did not significantly influence the TSS of the tomato, and fruit quality was not affected in various fertilizer treatments.

4. Conclusions

This study illustrates that combinations of mixed fertilizers or an integrated plant nutrient system (IPNS) produced the best results on plant growth, fruit measurements and yield of both varieties Roma VF and BARI 15. On the other hand, Roma VF produced higher quality attributes and yield over the BARI tomato 15. Combined applications of organic and inorganic sources of nutrients are more productive and sustain soil fertility. For the determination of an appropriate ratio of organic and inorganic sources, the experiment may be repeated at different locations for different varieties of tomato.

Acknowledgments: The authors express their sincere thanks to the USDA-FAS for providing funding support (DN) for this research and George Smith for reviewing manuscript.

Author Contributions: M. Ashraf Islam designed field experiment, conducted data analysis and prepared manuscript. Sumiya Islam assisted in the field data collection. Ayasha Akter helped in writing manuscript. M. Habibur Rahman helped in design and manuscript review. D. Nandwani overall supervised the project.

Conflicts of Interest: The authors declare no conflict of interest.

References

1. Aksoy, U. Ecological farming. II. In Proceedings of the Ecological Farming Symposium, Antalya, Turkey, 14–16 December 2001.
2. Chowdhury, R. Effects of chemical fertilizers on the surrounding environment and the alternative to the chemical fertilizers. *IES-ENVIS Newsl.* **2004**, *7*, 4–5.
3. Oyewole, C.I.; Opaluwa, H.; Omale, R. Response of Tomato (*Lycopersicon esculentum*): Growth and yield, to rates of mineral and poultry manure application in the guinea savanna agro-ecological zone in Nigeria. *J. Biol. Agric. Health Care* **2012**, *2*, 44–56.
4. Bulluck, L.R.; Ristaino, J.B. Effect of synthetic and organic soil fertility amendments on southern blight, soil microbial communities, and yield of processing tomatos. *Phytopathology* **2002**, *92*, 181–189. [[CrossRef](#)] [[PubMed](#)]
5. Bulluck, L.R.; Brosius, M.; Evanylo, G.K.; Ristaino, J.B. Organic and synthetic fertility amendments influence soil microbial, physical and chemical properties on organic and conventional farms. *Appl. Soil Ecol.* **2002**, *19*, 147–160. [[CrossRef](#)]
6. Arancon, N.Q.; Edwards, C.A.; Bierman, P.; Metzger, J.D.; Lee, S.; Welch, C. Effects of vermicompost on growth and marketable fruits of field-grown tomatoes, peppers and strawberries. *Pedobiologia* **2004**, *47*, 731–735. [[CrossRef](#)]
7. Heeb, A.; Lundegardh, B.; Savage, G.P.; Ericsson, T. Impact of organic and inorganic fertilizers on yield, taste, and nutritional quality of tomatoes. *J. Plant Nutr. Soil Sci.* **2006**, *169*, 535–541. [[CrossRef](#)]
8. Liu, B.; Gumpertz, M.L.; Hu, S.; Ristaino, J.B. Long-term effects of organic and synthetic soil fertility amendments on soil microbial communities and the development of southern blight. *Soil Biol. Biochem.* **2007**, *39*, 2302–2316. [[CrossRef](#)]
9. Maske, S.N.; Munde, G.R.; Maske, N.M. Effect of manures and fertilizer on brinjal (*Solanum melongena* L.) C.V. Krishna. *Bioinfolet* **2015**, *12*, 678–679.
10. Tonfack, L.B.; Bernadac, A.; Youmbi, E.; Mbouapouognigni, V.P.; Nguenguim, M.; Akoa, M. Impact of organic and inorganic fertilizers on tomato vigor, yield and fruit composition under tropical and soil conditions. *Fruits* **2009**, *64*, 167–177. [[CrossRef](#)]
11. Solaiman, A.R.M.; Rabbani, M.G. Effects of NPKS and cow dung on growth and yield of tomato. *Bull. Inst. Trop. Agric. Kyushu Univ.* **2006**, *1*, 31–37.
12. Haque, M.M.; Ilias, G.N.M.; Molla, A.H. Impact of *Trichoderma*-enriched biofertilizer on the growth and yield of mustard (*Brassica rapa* L.) and tomato (*Solanum lycopersicon* Mill.). *Agriculturists* **2012**, *10*, 109–119. [[CrossRef](#)]
13. Adesemoye, O.A.; Torbert, H.A.; Kloepper, J.W. Plant growth-promoting rhizobacteria allow reduced application rates of chemical fertilizers. *Microb. Ecol.* **2009**, *58*, 921–929. [[CrossRef](#)] [[PubMed](#)]
14. Sreenivasa, M.N.; Nagaraj, M.N.; Bhat, S.N. Beejamruth: A source for beneficial bacteria. *Karnataka J. Agric. Sci.* **2010**, *17*, 72–77.
15. Muhibullah, M.; Momotaz, S.; Chowdhury, A.T. Use of agrochemical fertilizers and their impact on soil, water and human health in the Khamargao Village of Mymensingh District, Bangladesh. *J. Agron.* **2005**, *4*, 109–115.
16. Food and Agriculture Organization (FAO). *Production Year Book*; Food and Agriculture Organization of the United Nations: Rome, Italy, 2010; Volume 53, pp. 132–133.
17. Bangladesh Bureau of Statistics (BBS). *BBS, Statistics Division, Ministry of Planning*; Govt. of People's Republic of Bangladesh: Dhaka, Bangladesh, 2003; pp. 1–7.
18. Gabal, M.R.; Abd-Allah, I.M.; Hass, F.M.; Hassannen, S. Evaluation of some American tomato cultivars grown for early summer production in Egypt. *Ann. Agric. Sci. Moshthor J.* **1984**, *22*, 487–500.

19. Nandwani, D. Growth and yield response of four tomato cultivars in the US Virgin Islands. *J. Agric. Univ. Puerto Rico* **2014**, *97*, 181–184.
20. Ahammad, K.U.; Siddiky, M.A.; Ali, Z.; Ahmed, R. Effects of planting time on the growth and yield of tomato varieties in late season. *Progress. Agric.* **2009**, *20*, 73–78. [[CrossRef](#)]
21. SAS Institute Inc. *SAS Technical Report R-109, Conjoint Analysis Examples*; SAS Institute Inc.: Cary, NC, USA, 1993.
22. Patil, M.B.; Mohammed, R.G.; Ghadge, P.M. Effect of organic and inorganic fertilizers on growth, yield and quality of tomato. *J. Maharashtra Agric. Univ.* **2004**, *29*, 124–127.
23. Reddy, C.S.; Narayanamma, M.; Chiranjeevi, C.; Reddy, L.P. Effect of nutrient sources on the fruit yield of tomato (*Lycopersicon esculentum* Mill.). *Veg. Sci.* **2002**, *29*, 193–194.
24. Golam, F. Effect of Spacing and Fertilizers on the Growth and Yield of Cabbage. Master's Thesis, Bangladesh Agricultural University, Mymensingh, Bangladesh, 2015.
25. Ullah, M.S.; Islam, M.S.; Islam, M.A.; Haque, T. Effects of organic manures and chemical fertilizers on the yield of brinjal and soil properties. *J. Bangladesh Agric. Univ.* **2008**, *6*, 271–276. [[CrossRef](#)]
26. Glick, B.R.; Todorovic, B.; Czarny, J.; Cheng, Z.; Duan, J.; McConkey, B. Promotion of plant growth by bacterial ACC deaminase. *Crit. Rev. Plant Sci.* **2007**, *26*, 227–242. [[CrossRef](#)]
27. Bona, E.; Cantamessa, S.; Massa, N.; Manassero, P.; Marsano, F.; Copetta, A.; Lingua, G.; D'Agostino, G.; Gamalero, E.; Berta, G. Arbuscular mycorrhizal fungi and plant growth-promoting pseudomonads improve yield, quality and nutritional value of tomato: A field study. *Mycorrhiza* **2017**, *27*, 1–11. [[CrossRef](#)] [[PubMed](#)]



© 2017 by the authors. Licensee MDPI, Basel, Switzerland. This article is an open access article distributed under the terms and conditions of the Creative Commons Attribution (CC BY) license (<http://creativecommons.org/licenses/by/4.0/>).